

Systematic review with meta-analysis: effect of fibre supplementation on chronic idiopathic constipation in adults

S. Christodoulides^{*,†}, E. Dimidi^{*,†}, K. C. Fragkos[‡], A. D. Farmer^{*,§}, K. Whelan[†] & S. M. Scott^{*}

^{*}The Wingate Institute for Neurogastroenterology and GI Physiology Unit, Centre for Neuroscience and Trauma, Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK.

[†]Diabetes and Nutritional Sciences Division, Faculty of Life Sciences and Medicine, King's College London, London, UK.

[‡]Centre for Gastroenterology and Clinical Nutrition, University College London, London, UK.

[§]University Hospitals of North Midlands, Stoke on Trent, UK.

Correspondence to:

Dr S. M. Scott, Wingate Institute, 26 Ashfield Street, Whitechapel, London E1 2AJ, UK.
E-mail: m.scott@qmul.ac.uk

Publication data

Submitted 14 October 2015
First decision 28 October 2015
Resubmitted 10 March 2016
Resubmitted 11 April 2016
Accepted 22 April 2016
EV Pub Online 12 May 2016

As part of AP&T's peer-review process, a technical check of this meta-analysis was performed by Dr Y. Yuan. This article was accepted for publication after full peer-review.

SUMMARY

Background

Chronic idiopathic constipation is a common symptom-based gastrointestinal disorder responsible for a substantial economic health service burden. Current guidelines recommend the use of fibre as a first-line treatment.

Aim

To investigate the effect of fibre (including prebiotic) supplementation on global symptom response, stool output, gut microbiota composition and adverse events in adults with chronic idiopathic constipation.

Methods

Medline, EmBase, Web of Science, Scopus and the Cochrane central register of controlled trials were searched through to February 2016. Conference proceedings from 2003 to 2015 were hand-searched. There were no language restrictions. Forest plots with 95% CIs were generated using a random-effects model.

Results

The search strategy generated 1072 citations, of which seven individual randomised controlled trials were eligible. Overall, 113 of 147 (77%) patients assigned to fibre responded to therapy, compared with 61 of 140 (44%) allocated to placebo (RR of success to respond 1.71, 95% CI 1.20–2.42, $P = 0.003$). Fibre significantly increased stool frequency (SMD, standardised mean difference = 0.39; 95% CI 0.03–0.76; $P = 0.03$) and softened stool consistency (SMD = 0.35; 95% CI 0.04–0.65; $P = 0.02$) compared with placebo. Flatulence was significantly higher with fibre compared to placebo (SMD 0.56, 0.12–1.00, $P = 0.01$). Overall quality of evidence was low.

Conclusions

This meta-analysis demonstrates that fibre is moderately effective, but also causes moderate gastrointestinal side effects. However, these findings need to be treated with caution due to a high risk of bias. Accordingly, further large, methodologically rigorous trials are required, before any definitive recommendation regarding its risk–benefit profile can be made.

PROSPERO registration number CRD42014007005.

Aliment Pharmacol Ther 2016; **44**: 103–116

INTRODUCTION

Chronic idiopathic constipation (CIC) is a heterogeneous symptom-based disorder with an estimated global prevalence of 14%¹ characterised by infrequent defecation, difficult stool passage or a combination of the two in the absence of an organic cause.² Females have been shown to be two to three times more likely to have CIC than men.³ CIC is associated with impaired quality of life,⁴ increased risk of colorectal cancer⁵ and is responsible for a substantial economic health service burden.⁶ The pathophysiology of CIC is multifactorial and incompletely understood. Beyond simple lifestyle advice (e.g. increasing fluid intake and levels of exercise), laxatives are a widely used treatment⁷ but are associated with sub-optimal outcomes due to variable efficacy, adverse events, cost, taste and inconvenience.⁸

First-line management as recommended in British, American, European and other global guidelines, as well as expert commentaries, is fibre supplementation.^{9–13} Fibre intake may accelerate whole gut transit time, by increasing luminal bulk resulting in increased peristalsis.¹⁴ Fibre can also influence bulking directly via water retention which also normalises stool form.^{15, 16} Furthermore, fermentation of fibre can increase stool bulk by increasing microbial biomass and fermentation by-products, such as short chain fatty acids.¹⁷ Gut transit time may also be indirectly accelerated through lowering of luminal pH and possibly through secondary effects on the gut microbiota.¹⁸ Nevertheless, up to 50% of patients do not respond or become dissatisfied with fibre as a treatment stratagem.¹⁹ The term fibre refers to carbohydrate polymers with three or more monomers that are not hydrolysed by endogenous enzymes in the human small intestine.²⁰ This definition also includes prebiotics, which are soluble fibres that are selectively fermented and result in specific changes in the composition and/or activity of the gut microbiota, and are postulated to confer health benefit(s) upon the host.²¹

The efficacy of fibre in the management of CIC in adults has been documented in a previous systematic review.²² However, to date there has been no definitive quantitative summary of available evidence and outcomes (particularly on the effect of prebiotics on gut microbiota composition). Thus, we aimed to address this knowledge gap by objectively assessing the effect of fibre supplementation on measures of: (i) response to therapy, (ii) stool output, (iii) gut microbiota composition, and (iv) adverse events in adults with CIC.

METHODS

A systematic review and meta-analysis was performed following the guidelines of the Cochrane Handbook for Systematic Reviews of Interventions,²³ and reported in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.²⁴

Search strategy and study selection

A literature search for randomised controlled trials (RCTs) evaluating fibre supplementation used in the treatment of CIC in adults was conducted. A database search was performed in February 2016 using MEDLINE, EMBASE, WEB OF SCIENCE, SCOPUS and The Cochrane Central Register of Controlled Trials. No date limitations were applied and searches were not restricted by language. Studies on CIC were identified with the terms: constipation OR gastrointestinal transit [both as medical subject headings (MeSH) and free text terms]. These were combined using the set operator AND with studies identified with the terms: fibre OR psyllium (both as MeSH terms and free text terms). Results were then further combined with the operator AND with highly sensitive search filter for identifying RCTs. The detailed search strategy for MEDLINE is presented in Table S1. Reference lists of eligible studies, as well as reference lists of previous systematic reviews on fibre and constipation, were manually scanned for additional studies not identified by the electronic searches. An attempt to identify completed but unpublished trials was performed by searching the ClinicalTrials.gov and www.isrctn.com databases. Finally, abstracts of the following conference proceedings were hand-searched to identify potentially relevant studies: Digestive Diseases Week (2003–2015), British Society of Gastroenterology Annual Meeting (2003–2015) and United European Gastroenterology Week (2009–2015). Titles and abstracts identified from the search strategy were evaluated by two independent investigators (SC and ED) using pre-defined eligibility criteria. The full text of any title or abstract deemed potentially eligible by either investigator was retrieved and foreign language papers were translated when required. Subsequently, the two reviewers independently assessed the eligibility of each full-text article and disagreements were resolved by consensus with a third researcher (SMS).

Eligibility criteria

The eligibility criteria were developed using the Population, Intervention, Comparator, Outcomes, Study

(PICOS) design approach.²⁵ *Population* – adults (≥ 18 years old) with CIC diagnosed using clinical diagnosis, self-report or Rome criteria. Only studies in which patients were recruited from community or out-patient settings were included. *Intervention* – studies using supplementary fibre as defined by the Commission of European Communities.²⁰ *Comparator* – studies comparing fibre (including prebiotic) supplements with placebo/control. Trials were also included if they reported interventions with supplementary fibre in combination with other ingredients (active comparators) as long as the effect of fibre could be isolated. *Outcomes* – studies reporting either dichotomous or continuous data on overall response to therapy, stool output, gut microbiota concentrations and adverse events. *Study design* – RCTs with parallel group design, or the first period of cross-over RCTs (to reduce the risk of carry-over effect) were considered eligible for inclusion.

Data extraction

Two investigators (SC and ED) independently extracted data on patient characteristics, interventions, comparators, measure outcomes and study design using standardised data extraction forms. Authors of included studies were contacted for missing data and their responses were included in the analyses. Whenever allowed by trial reporting, data were extracted as intention-to-treat analyses, with all drop-outs assumed to be treatment failures for dichotomous outcomes (i.e. no response to therapy). If this was not clear from the original article, an analysis on all patients with reported evaluable data was undertaken.

Assessment of risk of bias

The two investigators independently assessed risk of bias according to the Cochrane Collaboration handbook,²³ with disagreements resolved by discussion. Studies were assessed for the methods used to generate randomisation, conceal allocation, blinding, incomplete outcome data, selective reporting and other sources of bias. Furthermore, the overall quality of evidence (confidence in effect estimates) for each outcome was rated using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach.²⁶ Quality of evidence was rated from high to very low.²⁷

Data synthesis and statistical analysis

Data were pooled using a random-effects model to give a more conservative estimate of the effect of fibre supplementation on CIC, allowing for heterogeneity between

studies.²⁸ The estimates of treatment effects were expressed as relative risk (RR) and 95% confidence interval (CI) for dichotomous outcomes (response to therapy) and standardised mean difference (SMD) with 95% CI for continuous outcomes (stool frequency, stool consistency, gut microbiota concentrations and adverse events). SMD was used as a summary statistic for all continuous outcomes because the studies measured the same outcome using varied scales or because no details of the scale and scoring system were reported, which prevented the data being converted to the same unit, thus prevented calculation of the mean difference. SMD values of 0.2, 0.5 and 0.8 were defined as small, moderate and large effect size respectively.²⁹ The number needed to treat (NNT) and 95% CIs were calculated for response to therapy using the formula: $NNT = 1 / (1 - RR) \times \text{assumed control group risk}$.²³ Furthermore, the precision of the estimate of effect for the primary outcome was tested by calculating the optimal information size using α (0.05) and β (0.20) values, and a RR reduction of response to therapy of 30%.³⁰ Heterogeneity between studies was assessed using the I^2 statistic with a cut-off of 25%,³¹ and/or the chi-squared test, with a $P < 0.10$ to define significant heterogeneity. I^2 statistic values of 25%, 50% and 75% were defined as low, moderate and high heterogeneity respectively.³² Subgroup analyses were carried out to explore possible causes of heterogeneity and subgroup treatment interactions using the chi-squared significance test.³³ An interaction $P < 0.10$ was considered as a cut-off.³⁴ Forest plots with 95% CIs were generated for all outcomes using Review Manager v. 5.3 (Copenhagen, Denmark) and STATA v.12.0 (Stata Statistical Software; StataCorp, College Station, TX, USA). Funnel plots were generated, where sufficient number of studies were identified, to assess for evidence of asymmetry and therefore possible publication bias or other small study effects³⁵; these were evaluated by both observing the funnel plots and using Egger's test analysis.³⁶ A two-tailed $P \leq 0.05$ was adopted as the statistical criterion. The kappa statistic for inter-observer agreement between the two reviewers was also calculated.³⁷

RESULTS

Study selection

The search strategy generated 1072 citations, of which 1064 studies were identified from the primary electronic databases and eight studies identified through manual search. Of these, 377 were duplicates leaving 695 records to be screened, of which 52 were potentially relevant and

retrieved for full-text review (Figure 1). Of these, 45 were excluded with seven RCTs ultimately eligible.^{38–44} Agreement between investigators for trial eligibility was substantial (kappa statistic = 0.80).

Study characteristics

The seven RCTs comparing fibre supplementation with placebo/control involved a total of 430 adults with CIC.^{38–44} Six studies were published in English and one in Spanish.⁴² Authors of six of the trials were contacted to obtain supporting information about the methodology used.^{39–44} Of these, four provided responses that were included in the analyses.^{39–41, 44} Table 1 shows the definition of CIC used in the included studies, and Table 2 details the characteristics of the included studies. There was considerable variability in the type (e.g. psyllium, inulin) and dose (10–22.5 g/day) of fibre studied. Treatment periods varied from 2 to 8 weeks and the proportion of women in the trials ranged between 64% and 100%.

Efficacy and safety of fibre in the treatment of CIC

Response to therapy. Dichotomous data on response to therapy, measured by symptomatic improvement, were reported by four RCTs^{38, 41, 42, 44} including 287 patients. Overall, 113 of 147 (77%) patients assigned to fibre responded to therapy, compared with 61 of 140 (44%) allocated to placebo (RR of success to respond = 1.71; 95% CI 1.20–2.42; $P = 0.003$), with borderline heterogeneity between studies ($I^2 = 24%$, $\chi^2 P = 0.27$). Overall, the NNT with fibre to result in response in one patient was 3 (95% CI 2.6–3.4). Given the borderline heterogeneity, subgroup analyses were conducted (Table 3). The NNT were relatively stable in all these analyses. However, there was no heterogeneity between trials when only the two trials that used ‘proportion with no straining during defecation’ to define response to therapy ($I^2 = 0%$, $\chi^2 P = 0.77$), the two trials that used nonprebiotics ($I^2 = 0%$, $\chi^2 P = 0.69$), and when the three studies that used high dose ($I^2 = 0%$, $\chi^2 P = 0.61$) were included in the analysis. Subgroup analyses demonstrated no statistically significant differences in efficacy according to definition of response to therapy, type of fibre used, fibre solubility in water, and prebiotics vs. nonprebiotics (interaction $P > 0.10$). In contrast, the subgroup analysis evaluating low (≤ 15 g/day) vs. high dose (> 15 g/day) suggested an increased effect with high dose (interaction $P = 0.09$), partly explaining the heterogeneity observed across all studies. The optimal information size was calculated on the basis of a RR reduction of response to

therapy of 30%. Optimal information size (617 individuals) was greater than the total sample size (287 participants), while the number of events across trials was relatively low (174 events) (Table 4).

Stool frequency. Continuous data on stool frequency were reported by six studies including 406 patients.^{38–43} Overall, fibre supplementation significantly increased stool frequency compared with placebo (SMD = 0.39; 95% CI 0.03–0.76; $P = 0.03$), indicating a moderate effect size, albeit with statistically significant heterogeneity between studies ($I^2 = 56%$, $\chi^2 P = 0.04$) (Figure S1). Results of subgroup analyses (Figure 2) showed no differences in efficacy according to type of fibre used (interaction $P = 0.81$), and prebiotics vs. nonprebiotics (interaction $P = 0.91$), although psyllium and nonprebiotics increased stool frequency ($P = 0.0005$). In addition, there was no heterogeneity between trials when only the two trials that used psyllium, and the two trials that used nonprebiotics were included in the analysis ($I^2 = 0%$, $\chi^2 P = 0.87$). Subgroup analyses suggested a possible increased effect with high dose (interaction $P = 0.17$). High-dose (> 15 g/day) fibre supplementation significantly increased stool frequency compared with placebo (SMD = 0.77; 95% CI 0.03–1.51; $P = 0.04$), indicating a large effect size. However, statistically significant heterogeneity between studies existed ($I^2 = 71%$, $\chi^2 P = 0.06$). Results of subgroup analysis according to sample size revealed a borderline statistically significant increase in stool frequency when studies that recruited less than 50 participants ($P = 0.003$) were included in the analysis (interaction $P = 0.10$).

Stool consistency. Stool consistency was reported as a continuous end point in five studies including 346 patients.^{38–40, 42, 43} Overall, fibre supplementation significantly softened stool consistency compared with placebo (SMD = 0.35; 95% CI 0.04–0.65; $P = 0.02$), indicating a moderate effect size, however heterogeneity between studies existed ($I^2 = 34%$, $\chi^2 P = 0.19$) (Figure S2). Heterogeneity between trials was lower when only the two trials that used psyllium and when only the two trials that used nonprebiotics were included in the analysis ($I^2 = 8%$, $\chi^2 P = 0.30$). Subgroup analyses showed no difference in efficacy according to type of fibre used (interaction $P = 0.18$) (Figure 3). However, psyllium ($P = 0.009$) and a mixture of inulin with resistant maltodextrin ($P = 0.008$) softened stool consistency. Results of the subgroup analysis comparing prebiotics vs. nonprebiotics showed




Figure 1 | PRISMA flow diagram of studies in systematic review.

borderline statistically significant improvement in stool consistency when studies that used nonprebiotics ($P = 0.009$) or a mixture of a prebiotic/nonprebiotic

($P = 0.008$) were included in the analysis (interaction $P = 0.10$). The subgroup hypothesis evaluating single fibres vs. mixtures of (multiple) fibre suggested an

Table 1 | Definition of chronic idiopathic constipation used in randomised controlled trials of fibre supplementation in adults

Study	Diagnostic criteria for CIC	Details of diagnostic criteria
Fenn, 1986 ³⁸	Clinical diagnosis	Patients suffering from functional constipation
Ashraf, 1995 ³⁹	Clinical diagnosis	Patients with chronic idiopathic constipation confirmed by prospectively administered stool diaries
Marteau, 2011 ⁴⁰	Rome III criteria	Patients suffering from functional constipation according to the Rome definition (<3 stools/week and/or straining in defecation)
Waitzberg, 2012 ⁴¹	Self-reported	Females with at least 3 months of primary constipation defined as less than three bowel movements per week
Lopez Roman, 2008 ⁴²	Rome II criteria	Patients meeting the Rome II criteria for chronic idiopathic constipation
Surakka, 2009 ⁴³	Self-reported	Patients with difficulties in intestinal function (including fewer than five BM per week or continuous difficulties in defecation or both)
Badiali, 1995 ⁴⁴	Clinical diagnosis	Patients with prolonged large bowel transit seeking medical advice for chronic primary constipation

CIC, chronic idiopathic constipation; BM, bowel movements.

Table 2 | Characteristics of included randomised controlled trials of fibre supplementation in chronic idiopathic constipation in adults

Study	Country	Setting	Study design	Criteria used to define response to therapy	No. of patients (% female)	Age Mean (range) (years)	Fibre used	Water solubility	Prebiotics	Daily Dose (g)	Form	Duration of therapy
Fenn, 1986 ³⁸	UK	Primary care	Single blind, parallel group	Proportion with an improvement in global symptoms	201 (75)	49 (17–70)	Psyllium	Soluble	Nonprebiotic	19.2	Powder	2 weeks
Ashraf, 1995 ³⁹	USA	Tertiary care	Double blind, parallel group	NR	22 (64)	51 (40–75)	Psyllium	Soluble	Nonprebiotic	10.2	Powder	8 weeks
Marteau, 2011 ⁴⁰	France	Primary care	Double blind, parallel group	NR	50 (88)	57 (50–70)	Inulin	Soluble	Prebiotic	15	Powder	4 weeks
Waitzberg, 2012 ⁴¹	Brazil	Primary care	Double blind, parallel group	Proportion with constipation relief	60 (100)	38 (18–65)	Inulin and PHGG	Soluble	Prebiotic and nonprebiotic	15	Powder	3 weeks
Lopez Roman, 2008 ⁴²	Spain	Tertiary care	Double blind, parallel group	Proportion with no straining during defecation	32 (88)	47 (17–77)	Inulin and RM	Soluble	Prebiotic and nonprebiotic	22.5	Milk	20 days
Surakka, 2009 ⁴³	Finland	Tertiary care	Double blind, crossover	NR	41 (76)	68 (60–80)	GOS	Soluble	Prebiotic	10	Yoghurt	3 weeks
Badiali, 1995 ⁴⁴	Italy	Tertiary care	Double blind, crossover	Proportion with no straining during defecation	24 (92)	41 (18–65)	Wheat bran	Insoluble	Nonprebiotic	19.8	Powder	4 weeks

GOS, galacto-oligosaccharides; NR, not reported (study did not report any dichotomous data on response to therapy); PHGG, partially hydrolysed guar gum; RM, resistant maltodextrin.

increased effect with mixtures of fibre ($P = 0.008$) (interaction $P = 0.09$). Furthermore, there were subgroup differences by dose of fibre used (interaction $P = 0.06$)

showing an increased effect on stool consistency with high dose (>15 g/day) ($P = 0.006$), partly explaining the heterogeneity observed across all studies.

Table 3 | Subgroup analyses of efficacy of fibre resulting in response to therapy in adults with chronic idiopathic constipation

	Number of studies	Number of patients	RR of success to respond to therapy (95% CI)	Sig.	Interaction P value	χ^2 (P value)	I^2 (%)	Number needed to treat (95% CI)
All studies	4 ^{38, 41, 42, 44}	287	1.71 (1.2–2.42)	0.003*		0.27	24	3 (2.6–3.4)
Definition of response to therapy					0.23			
Proportion with no straining during defecation	2 ^{42, 44}	44	2.70 (1.19–6.11)	0.02*		0.77	0	2 (1.7–2.3)
Other definition	2 ^{38, 41}	243	1.46 (0.82–2.58)	0.20		0.10	64	N/A
Type of fibre					0.90			
Psyllium	1 ³⁸	197	1.80 (1.45–2.25)	<0.00001*		N/A	N/A	3 (2.9–3.1)
Wheat bran	1 ⁴⁴	16	2.33 (0.66–8.22)	0.19		N/A	N/A	N/A
Mixture of inulin with PHGG or RM	2 ^{41, 42}	74	1.59 (0.53–4.75)	0.41		0.08*	67	N/A
Fibre solubility					0.61			
Soluble fibre	3 ^{38, 41, 42}	271	1.65 (1.03–2.62)	0.04*		0.16	46	3 (2.5–3.5)
Insoluble fibre	1 ⁴⁴	16	2.33 (0.66–8.22)	0.19		N/A	N/A	N/A
Prebiotics vs. nonprebiotics					0.81			
Mixture of a prebiotic/nonprebiotic	2 ^{41, 42}	74	1.59 (0.53–4.75)	0.41		0.08*	67	N/A
Nonprebiotics	2 ^{38, 44}	213	1.82 (1.46–2.26)	<0.00001*		0.69	0	2 (1.9–2.1)
Dose of treatment					0.09*			
Low dose (≤ 15 g)	1 ⁴¹	46	0.98 (0.49–1.96)	0.96		N/A	N/A	N/A
High dose (>15 g)	3 ^{38, 42, 44}	241	1.85 (1.50–2.29)	<0.00001*		0.61	0	2 (1.8–2.2)

CI, confidence interval; interaction P value, P value of tests for interactions; N/A, not applicable (too few studies to assess heterogeneity and/or relative risk not statistically significant to calculate number needed to treat); PHGG, partially hydrolysed guar gum; RM, resistant maltodextrin; RR, relative risk; Sig., significance.

* Statistically significant.

Gut microbiota concentrations. The effect of fibre supplementation on faecal bifidobacteria and clostridia concentrations was reported in three studies. Bacterial counts were assessed via molecular techniques (two studies used the quantitative polymerase chain reaction^{40, 41} and one study used fluorescence *in situ* hybridisation.⁴³) Bifidobacteria counts were reported in three studies, including 151 patients.^{40, 41, 43} Overall, fibre supplementation did not significantly increase bifidobacteria numbers compared with placebo (SMD = 0.43; 95% CI –0.20 to 1.07; $P = 0.18$), and statistically significant heterogeneity between studies existed ($I^2 = 72\%$, $\chi^2 P = 0.03$) (Figure S3). Subgroup analyses were performed to explore possible causes of the significant heterogeneity observed (Figure 4). The subgroup analysis comparing prebiotics vs. nonprebiotics showed a statistically significant increase in bifidobacteria counts with prebiotics (SMD = 0.75; 95% CI 0.33–1.18; $P = 0.0005$) [inulin ($P = 0.02$) and galacto-oligosaccharides (GOS)

($P = 0.01$)] (interaction $P = 0.008$). In addition, there was no heterogeneity between trials when only the two trials that used prebiotics were included in the analysis ($I^2 = 0\%$, $\chi^2 P = 0.78$) (Figure S3). Two studies, including 101 patients, reported data on clostridia counts.^{41, 43} Clostridia levels were significantly reduced with fibre supplementation compared to placebo (SMD = –0.66; 95% CI –1.29 to –0.02; $P = 0.04$), indicating a moderate effect size, although heterogeneity between studies existed ($I^2 = 56\%$, $\chi^2 P = 0.13$).

Adverse events. None of the studies reported total number of adverse events or patient withdrawal because of fibre side effects. Nevertheless, three studies reported continuous data regarding individual gastrointestinal adverse events (flatulence and bloating). Three studies, including 115 patients, provided data on flatulence.^{40, 43, 44} Overall, flatulence was significantly higher with fibre supplementation compared to placebo

Table 4 | GRADE system approach for quality of evidence assessment of each outcome

Outcome	No. of studies	No. of patients	Quality assessment					Publication bias	Quality of evidence
			Study limitations	Inconsistency	Indirectness	Imprecision			
Response to therapy	4 ^{38, 41, 42, 44}	287	Serious limitations*	⊕	⊕	OIS greater than number of patients; relatively low number of events	Not evaluable	⊕⊕⊕⊕, Low	
Stool frequency	6 ^{38–43}	406	Serious limitations*	Significant heterogeneity between studies; variability in results	⊕	Modest number of patients studied with wide 95% CIs	⊕	⊕⊕⊕⊕⊕, Low	
Stool consistency	5 ^{38–40, 42, 43}	346	Serious limitations*	Moderate heterogeneity between studies	⊕	Very modest number of patients studied with wide 95% CIs	Not evaluable	⊕⊕⊕⊕, Very low	
Gut microbiota concentrations	3 ^{40, 41, 43}	151	Serious limitations*	Significant heterogeneity between studies; variability in results	⊕	Only a small number of patients studied with wide 95% CIs	Not evaluable	⊕⊕⊕⊕, Very low	
Adverse events	3 ^{40, 43, 44}	115	Serious limitations*	⊕	⊕	Only a small number of patients studied with wide 95% CIs	Not evaluable	⊕⊕⊕⊕, Low	

CI, confidence interval; OIS, optimal information size.

* All trials were at high risk of bias and all failed to adhere to an intention-to-treat analysis; there was also high attrition (incomplete outcome data) and reporting bias (selective reporting) among trials; ⊕, the criterion was fulfilled; ⊖, the criterion was not fulfilled.

(SMD = 0.56; 95% CI 0.12–1.00; $P = 0.01$), indicating a moderate effect size, with low heterogeneity between studies detected ($I^2 = 23\%$, $\chi^2 P = 0.27$) (Figure S4). Bloating was reported in three studies, including 115 patients.^{40, 43, 44} Fibre supplementation increased bloating compared with placebo (SMD = 0.36; 95% CI -0.01 to 0.74; $P = 0.06$) with borderline statistical significance. No heterogeneity between studies existed ($I^2 = 0\%$, $\chi^2 P = 0.94$) (Figure S5).

Publication bias

The studies identified herein were too few in number to assess for evidence of asymmetry, and in turn for evidence of publication bias or other small study effects.³⁶

Risk of bias

None of the studies was at low risk of bias, both at the study (Figure S6) and outcome level (data available on request), and none of the trials followed an intention-to-treat analysis. Attrition bias (incomplete outcome data) and reporting bias (selective reporting) were prevalent, whereas performance bias (blinding of participants and personnel) and other bias were low among trials, both at

the study (Figure S6) and outcome level (data available on request). GRADE criteria were applied to interpret results and assess overall quality of evidence (confidence in effect estimates) for each outcome (Table 4). The quality of evidence in terms of response to therapy, stool frequency, and adverse events was graded as low, whereas the confidence in the effect estimates for stool consistency and gut microbiota concentrations was graded as very low.

DISCUSSION

This meta-analysis demonstrates that fibre supplementation is more effective than placebo for the treatment of CIC in adults. However, there is only low-quality evidence that fibre is moderately effective. Furthermore, individual gastrointestinal side effects (e.g. flatulence) were higher in patients receiving fibre, although no patient withdrew because of fibre side effects. Only four trials^{38, 41, 42, 44} adhered to the ROME committee's recommendations⁴⁵ by reporting global dichotomous data on response to therapy. The NNT with fibre to result in response to therapy was between 2 and 3, with subgroup analyses suggesting that high dose (>15 g) was the most




Figure 2 | Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for stool frequency among subgroups with the use of a random-effects model. GOS, galacto-oligosaccharides; PHGG, partially hydrolysed guar gum; RM, resistant maltodextrin.

effective, with a NNT of 2. However, the definitions of response to therapy were inconsistent, and placebo rate of 44% was very high (e.g. in comparison to a placebo response of 27%, shown in a meta-analysis of laxatives in CIC).⁴⁶ In addition, the optimal information size (617 individuals) was not met (287 participants), thus lowering confidence in estimates of effect for imprecision.³⁰ Hence, the effect size and NNT obtained from this meta-analysis need to be treated with caution.

Prebiotics (inulin and GOS) seem to have no benefit over placebo in increasing stool frequency. The most robust evidence for an individual fibre seems to be for psyllium (a nonprebiotic). High fibre dose (> 15 g) was found to increase stool frequency with a large effect size. The most recent meta-analysis on the efficacy of laxatives in CIC demonstrated similar efficacy to the data for high fibre dose⁴⁶; both osmotic laxatives, including

polyethylene glycol (~ 17 g once or twice daily), and stimulant laxatives [e.g. bisacodyl (10 mg)] significantly increased stool frequency with a large effect size.⁴⁶ However, the fact that small sample-sized studies (< 50 participants) showed higher increases in stool frequency suggests the possibility of small study bias, given that smaller studies tend to show larger estimated effects than larger studies.⁴⁷

Fibre supplementation moderately softened stool consistency, also with a dose-dependent effect. Psyllium showed greater effect on stool consistency compared to prebiotics (GOS and inulin). Psyllium (ispaghula husk), obtained from the seeds of *Plantago ovata*, is a soluble viscous fibre with a high water-holding capacity that normalises stool form.⁴⁸ Stool water content has been highly associated with stool consistency, and a relatively small increase in stool water content has been shown to




Figure 3 | Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for stool consistency among subgroups with the use of a random-effects model. GOS, galacto-oligosaccharides; RM, resistant maltodextrin.

result in a relatively large stool softening effect.^{15, 49} GOS and inulin are highly fermentable and are almost completely fermented in the colon, and despite high initial water-holding capacity, their fermentation results in a loss of water-holding capacity and thus they have little effect on stool consistency.¹⁶ In contrast, psyllium is only partially fermented in the gut and the stool softening effect is a direct consequence of its ability to form a gel and hold many times its own weight in water.¹⁶

In terms of side effects, flatulence and bloating were moderately increased after fibre consumption with a trend towards a dose-dependent effect. This might partly explain why patients (especially those with pre-existing symptoms of bloating) often poorly tolerate fibre supplementation.¹⁸ However, some degree of spontaneous unblinding was arguably present in these trials because of the effect of fibre on bowel symptoms, and it is likely to have affected subjective symptom assessments.⁵⁰

Preliminary data have proposed a role for abnormal gut microbiota composition, so-called dysbiosis, in the pathogenesis of CIC.⁵¹ Patients with CIC have been found to have lower concentrations of 'beneficial' bacteria (e.g. bifidobacteria) and higher concentrations of 'pathogenic' bacteria (e.g. clostridia).⁵¹ Pooled results with prebiotics (GOS and inulin) showed a statistically significant increase in bifidobacteria counts with a large effect size. Both GOS and inulin can be metabolised by bifidobacteria,⁵² though GOS is more bifidogenic,⁵³ compatible with a higher effect size. In addition, fibre supplementation significantly lowered clostridia counts compared to placebo, with a moderate effect size. However, whether these beneficial quantitative changes in gut microbiota composition can lead to changes in gastrointestinal transit, which may benefit symptoms of constipation, is still unknown. Consequently, findings in relation to the microbiota-dependent or microbiota-independent




Figure 4 | Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for faecal bifidobacteria concentrations among subgroups with the use of a random-effects model. GOS, galacto-oligosaccharides; PHGG, partially hydrolysed guar gum.

effect of fibre on gastrointestinal transit require confirmation in future studies.

Strengths and limitations

This systematic review and meta-analysis strictly adhered to recommended methodology, including a comprehensive search strategy, which included searching the 'grey' literature to minimise publication bias. Only RCTs were selected, and foreign language articles were translated to further minimise bias. Both the eligibility assessment and data extraction were performed independently and in duplicate. Authors of individual studies were also contacted in order to obtain supporting information, where required. We used an intention-to-treat analysis and data were pooled with a random-effects model to provide a more conservative estimate of the efficacy of fibre in CIC. We also conducted subgroup analyses to assess heterogeneity in results. Finally, we used the GRADE approach, an internationally recognised method of rating evidence, to assess quality of evidence in addition to reporting risk of bias. Limitations of the present study derive from the quality and reporting of the included trials. None of the studies identified was at low risk of bias, both at the study and outcome level, sufficient in itself to negatively affect the interpretation of results.⁵⁴ Furthermore, none of the trials followed an intention-to-treat analysis potentially resulting in an over-estimation of efficacy of fibre supplementation.⁵⁵ Application of

GRADE criteria indicated results were of low quality, mainly due to study limitations, inconsistency and imprecision. Controversy remains over whether a meta-analysis should be performed due to concerns over the methodological quality of the studies identified, due to the considerable variability in the type of fibre studied and due to the different habitual fibre intake level among the participants. The detection of heterogeneity in some of our analyses might reflect this variability. Different types of fibre have different physiological properties,⁵⁰ and therefore, the therapeutic benefit may be fibre-specific. Unfortunately, the small number of studies available with each type of fibre in the subgroup analyses did not enable us to perform robust type-specific comparisons, and thus results should be interpreted with caution. Insoluble fibre has been also shown to alter colonic function through a stool bulking effect¹⁷ and through mechanical stimulation/irritation of the gut mucosa,^{56, 57} but only one trial used insoluble fibre.⁴⁴

Comparison with other studies

To the best of our knowledge, this is the first meta-analysis to assess available data on the efficacy of fibre supplementation in adults with CIC. Another recent meta-analysis⁵⁸ mainly focused on paediatric patients, with only one trial including adult patients.⁴⁴ This study also included patients with organic constipation rather than being conducted exclusively in patients with CIC. A

systematic review performed exclusively in adult patients with CIC was published in 2011,²² and comprised six trials.^{38, 39, 42, 44, 59, 60} Of these, we excluded two studies^{59, 60} (in which the effect of fibre supplementation could not be isolated), and identified three additional studies,^{40, 41, 43} two of which^{40, 41} were published in the intervening 4 years, underlining the continuing interest in the manipulation of the gut microbiota through prebiotic supplementation as a potential therapy for CIC. Our meta-analysis demonstrated that prebiotic supplementation can beneficially alter gut microbiota concentrations. Within the most recent American College of Gastroenterology guidelines on the management of CIC in adults,¹¹ a meta-analysis examining the effect of fibre on response to therapy identified only three trials,^{38, 42, 59} with pooled results suggesting a beneficial effect of fibre compared to placebo, with a NNT of 2, similar to that found in the current study. Finally, a systematic review without meta-analysis by Rao *et al.* was published in 2015,⁶¹ though the database search for this review was limited to a 10-year time period (2004–2014). Furthermore, studies with crossover design and short washout periods (1–2 weeks) were considered eligible, thus increasing the carry-over effect risk. Moreover, not all studies included were RCTs, and studies that used natural fibre (e.g. prunes), instead of a fibre supplement, were not excluded. Prunes also contain sorbitol, a polyol with osmotic laxative effect,⁶² which might have skewed results.

Implications for fibre supplementation recommendations

Fibre supplements are food products and therefore are relatively safe, inexpensive and are widely available.⁶³ Moreover, their use has been also associated with a diminution in colorectal cancer risk compared to nonfibre laxative use.⁶⁴ Therefore, they are a reasonable first-line therapy for CIC albeit with the caveat that up to 50% of patients will be dissatisfied with their effect.¹⁹ It remains unclear which type of fibre is the most efficacious, although the most robust evidence seems to be for psyllium (a soluble, nonprebiotic fibre). It also remains unclear whether one type of fibre will provide relief to all patients, or whether there may be particular subtypes of this heterogeneous condition that might respond preferentially.⁶⁵ The aforementioned dissatisfaction with fibre is likely to be a consequence of increased gastrointestinal adverse events, such as flatulence and bloating,⁸ and warrants discussion with the patient prior to institution.

CONCLUSION

Current guidelines recommend the use of fibre supplementation as first-line treatment for CIC. However, overall there is only low-quality evidence to support such a recommendation. Meta-analysis demonstrated that fibre supplementation is moderately effective with a dose-dependent effect and also causes moderate gastrointestinal side effects, again in a dose-dependent manner. Nevertheless, none of the studies included were at low risk of bias, sufficient in itself to negatively affect the interpretation of results. Hence, the paucity of high-quality data highlights the need for further large, methodologically rigorous RCTs of fibre supplementation in CIC, adhering to the ROME Foundation criteria, before any recommendation regarding its risk–benefit profile can be definitively promoted.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Figure S1. Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for stool frequency with the use of a random-effects model. IV, inverse variance.

Figure S2. Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for stool consistency with the use of a random-effects model. IV, inverse variance.

Figure S3. Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for faecal bifidobacteria concentrations (prebiotics vs. nonprebiotics) with the use of a random-effects model. IV, inverse variance.

Figure S4. Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for flatulence (low dose vs. high dose) with the use of a random-effects model. IV, inverse variance.

Figure S5. Forest plot of randomised controlled trials in adults with chronic idiopathic constipation comparing fibre with placebo/control. Standardised mean differences (95% CIs) for bloating (low dose vs. high dose) with the use of a random-effects model. IV, inverse variance.

Figure S6. Overall risk of bias summary and risk of bias graph for all studies and outcomes together (study level).

Table S1. Detailed search strategy – Medline.

AUTHORSHIP

Guarantor of the article: SMS.

Author contributions: SC, SMS and KW conceived and designed the review. SC, SMS and KW developed the review protocol. SC and ED performed eligibility screening, carried out the data extraction and methodological quality assessment. SMS provided advice and arbitration on the selection process, data extraction and methodological quality assessment. SC and KCF analysed the data. SC, ADF, SMS and KW interpreted the data. SC wrote the original draft, and the other authors revised the draft critically for important intellectual

content and approved the final version of the paper, including the authorship list.

ACKNOWLEDGEMENTS

We thank Alicia Green, Chung Lee, Anke Nijhuis, Ausra Zdanaviciene and Anicha Nazim Ahmad for their assistance with the translation of foreign language articles. We also thank Eamonn Quigley, Cathy Signoret, Dan Waitzberg and Danilo Badiali for providing extra information on their studies.

Declaration of personal and funding interests: None.

REFERENCES

- Suares NC, Ford AC. Prevalence of, and risk factors for, chronic idiopathic constipation in the community: systematic review and meta-analysis. *Am J Gastroenterol* 2011; **106**: 1582–91.
- Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006; **130**: 1480–91.
- McCrea GL, Miaskowski C, Stotts NA, Macera L, Paul SM, Varma MG. Gender differences in self-reported constipation characteristics, symptoms, and bowel and dietary habits among patients attending a specialty clinic for constipation. *Gen Med* 2009; **6**: 259–71.
- Belsey J, Greenfield S, Candy D, Geraint M. Systematic review: impact of constipation on quality of life in adults and children. *Aliment Pharmacol Ther* 2010; **31**: 938–49.
- Guerin A, Mody R, Fok B, *et al.* Risk of developing colorectal cancer and benign colorectal neoplasm in patients with chronic constipation. *Aliment Pharmacol Ther* 2014; **40**: 83–92.
- Choung RS, Branda ME, Chitkara D, *et al.* Longitudinal direct medical costs associated with constipation in women. *Aliment Pharmacol Ther* 2011; **33**: 251–60.
- Wald A, Scarpignato C, Mueller-Lissner S, *et al.* A multinational survey of prevalence and patterns of laxative use among adults with self-defined constipation. *Aliment Pharmacol Ther* 2008; **28**: 917–30.
- Müller-Lissner S, Tack J, Feng Y, Schenck F, Specht Gryp R. Levels of satisfaction with current chronic constipation treatment options in Europe – an internet survey. *Aliment Pharmacol Ther* 2013; **37**: 137–45.
- National Institute for Health and Care Excellence. Constipation in children and young people. *NICE Qual Stand* 2014; **62**: 1–38.
- Bharucha AE, Pemberton JH, Locke GR 3rd. American Gastroenterological Association technical review on constipation. *Gastroenterology* 2013; **144**: 218–38.
- Ford AC, Moayyedi P, Lacy BE, *et al.* American College of Gastroenterology monograph on the management of irritable bowel syndrome and chronic idiopathic constipation. *Am J Gastroenterol* 2014; **109**: S2–26.
- Tack J, Muller-Lissner S, Stanghellini V, *et al.* Diagnosis and treatment of chronic constipation – a European perspective. *Neurogastroenterol Motil* 2011; **23**: 697–710.
- World Gastroenterology Organization. Practice guidelines. *Constipation* 2007; **1**: 1–10.
- Brownlee IA. The physiological roles of dietary fibre. *Food Hydrocolloids* 2011; **25**: 238–50.
- McRorie J, Pepple S, Rudolph C. Effects of fiber laxatives and calcium docusate on regional water content and viscosity of digesta in the large intestine of the pig. *Dig Dis Sci* 1998; **43**: 738–45.
- Blackwood AD, Salter J, Dettmar PW, Chaplin MF. Dietary fibre, physicochemical properties and their relationship to health. *J R Soc Promot Health* 2000; **120**: 242–7.
- Stephen AM, Cummings JH. Mechanism of action of dietary fibre in the human colon. *Nature* 1980; **284**: 283–4.
- Eswaran S, Muir J, Chey WD. Fiber and functional gastrointestinal disorders. *Am J Gastroenterol* 2013; **108**: 718–27.
- Johanson JF, Kralstein J. Chronic constipation: a survey of the patient perspective. *Aliment Pharmacol Ther* 2007; **25**: 599–608.
- Commission of European Communities. Commission Directive 2008/100/EC amending Council Directive 90/496/EEC on nutrition labelling for foodstuffs as regards recommended daily allowances, energy conversion factors and definitions. *Off J Eur Union* 2008; **285**: 9–13.
- Gibson GR, Scott KP, Rastall RA, *et al.* Dietary prebiotics: current status and new definition. *Food Sci Technol Bull* 2010; **7**: 1–19.
- Suares NC, Ford AC. Systematic review: the effects of fibre in the management of chronic idiopathic constipation. *Aliment Pharmacol Ther* 2011; **33**: 895–901.
- Higgins JPT, Green S. Cochrane Handbook for systematic reviews of interventions. The Cochrane Collaboration 2011; version 5.1.0. Available at: www.cochrane-handbook.org (accessed 15 September 2015).
- Liberati A, Altman DG, Tetzlaff J, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009; **339**: 1–27.
- Moher D, Tricco AC. Issues related to the conduct of systematic reviews: a focus on the nutrition field. *Am J Clin Nutr* 2008; **88**: 1191–9.
- Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schunemann HJ. What is “quality of evidence” and why is it important to clinicians? *BMJ* 2008; **336**: 995–8.
- Balshem H, Helfand M, Schunemann HJ, *et al.* GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011; **64**: 401–6.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**: 177–88.
- Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale, NJ: Lawrence Erlbaum Associates, 1988.
- Guyatt GH, Oxman AD, Kunz R, *et al.* GRADE guidelines 6. Rating the quality of evidence – imprecision. *J Clin Epidemiol* 2011; **64**: 1283–93.
- Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**: 1539–58.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; **327**: 557–60.

33. Altman DG, Bland JM. Interaction revisited: the difference between two estimates. *BMJ* 2003; **326**: 219–9.
34. Sun X, Briel M, Walter SD, Guyatt GH. Is a subgroup effect believable? Updating criteria to evaluate the credibility of subgroup analyses. *BMJ* 2010; **340**: 850–4.
35. Egger M, Davey-Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629–34.
36. Sterne JA, Sutton AJ, Ioannidis JP, *et al.* Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomized controlled trials. *BMJ* 2011; **342**: 1–8.
37. Sands ML, Murphy JR. Use of kappa statistic in determining validity of quality filtering for meta-analysis: a case study of the health effects of electromagnetic radiation. *J Clin Epidemiol* 1996; **49**: 1045–51.
38. Fenn GC, Wilkinson PD, Lee CE, Akbar FA. A general practice study of the efficacy of Regulan in functional constipation. *Br J Clin Pract* 1986; **40**: 192–7.
39. Ashraf W, Park F, Lof J, Quigley EMM. Effects of psyllium therapy on stool characteristics, colon transit and anorectal function in chronic idiopathic constipation. *Aliment Pharmacol Ther* 1995; **9**: 639–47.
40. Marteau P, Jacobs H, Cazaubiel M, Signoret C, Prevel JM, Housez B. Effects of chicory inulin in constipated elderly people: a double-blind controlled trial. *Int J Food Sci Nutr* 2011; **62**: 164–70.
41. Waitzberg DL, Pereira CCA, Logullo L, *et al.* Microbiota benefits after inulin and partially hydrolyzed guar gum supplementation – a randomized clinical trial in constipated women. *Nutr Hosp* 2012; **27**: 123–9.
42. López Román J, Martínez González AB, Luque A, *et al.* The effect of a fibre enriched dietary milk product in chronic primary idiopathic constipation. *Nutr Hosp* 2008; **23**: 12–9.
43. Surakka A, Kajander K, Rajilic M, *et al.* Yoghurt containing galactooligosaccharides facilitates defecation among elderly subjects and selectively increases the number of bifidobacteria. *Int J Probiotics Prebiotics* 2009; **4**: 65–74.
44. Badiali D, Corazziari E, Habib FI, *et al.* Effect of wheat bran in treatment of chronic nonorganic constipation. A double-blind controlled trial. *Dig Dis Sci* 1995; **40**: 349–56.
45. Irvine EJ, Whitehead WE, Chey WD, *et al.* Design of treatment trials for functional gastrointestinal disorders. *Gastroenterology* 2006; **130**: 1538–51.
46. Ford AC, Suares NC. Effect of laxatives and pharmacological therapies in chronic idiopathic constipation: systematic review and meta-analysis. *Gut* 2011; **60**: 209–18.
47. Harbord RM, Egger M, Sterne JA. A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Stat Med* 2006; **25**: 3443–57.
48. Marlett JA, Fischer MH. The active fraction of psyllium seed husk. *Proc Nutr Soc* 2003; **62**: 207–9.
49. Wenzl HH, Fine KD, Schiller LR, Fordtran JS. Determinants of decreased fecal consistency in patients with diarrhoea. *Gastroenterology* 1995; **108**: 1729–38.
50. Wood L, Egger M, Gluud LL, *et al.* Empirical evidence of bias in treatment effect estimates in controlled trials with different interventions and outcomes: meta-epidemiological study. *BMJ* 2008; **336**: 601–5.
51. Khalif IL, Quigley EMM, Konovitch EA, Maximova ID. Alterations in the colonic flora and intestinal permeability and evidence of immune activation in chronic constipation. *Dig Liver Dis* 2005; **37**: 838–49.
52. Falony G, Lazidou K, Verschaeren A, Weck S, Maes D, De Vuyst L. In vitro kinetic analysis of fermentation of prebiotic inulin-type fructans by *bifidobacterium* species reveals four different phenotypes. *Appl Environ Microbiol* 2009; **75**: 454–61.
53. Maathuis AJH, van den Heuvel EG, Schoterman MHC, Venema K. Galactooligosaccharides have prebiotic activity in a dynamic in vitro colon model using a ¹³C-labeling technique. *J Nutr* 2012; **142**: 1205–12.
54. Higgins JPT, Altman DG, Gotzsche PC, *et al.* The Cochrane Collaboration's tool for assessing risk of bias in randomized trials. *BMJ* 2011; **343**: 1–9.
55. Juni P, Altman DG, Egger M. Assessing the quality of controlled clinical trials. *BMJ* 2001; **323**: 42–6.
56. Tomlin J, Read NW. Laxative properties of indigestible plastic particles. *BMJ* 1988; **297**: 1175–6.
57. McIntyre A, Vincent RM, Perkins AC, Spiller RC. Effect of bran, ispaghula, and inert plastic particles on gastric emptying and small bowel transit in humans: the role of physical factors. *Gut* 1997; **40**: 223–7.
58. Yang J, Wang H, Zhou L, Xu C. Effect of dietary fiber on constipation: a meta analysis. *World J Gastroenterol* 2012; **18**: 7378–83.
59. Odes HS, Madar Z. A double-blind trial of a celandin, aloe vera and psyllium laxative preparation in adult patients with constipation. *Digestion* 1991; **49**: 65–71.
60. Hongisto S, Pajanen L, Saxelin M, Korpela R. A combination of fibre-rich rye bread and yoghurt containing *Lactobacillus GG* improves bowel function in women with self-reported constipation. *Eur J Clin Nutr* 2006; **60**: 319–24.
61. Rao SSC, Yu S, Fedewa A. Systematic review: dietary fibre and FODMAP-restricted diet in the management of constipation and irritable bowel syndrome. *Aliment Pharmacol Ther* 2015; **41**: 1256–70.
62. Attaluri A, Donahoe R, Valestin J, Brown K, Rao SSC. Randomised clinical trial: dried plums (prunes) vs. psyllium for constipation. *Aliment Pharmacol Ther* 2011; **33**: 822–8.
63. Schmier JK, Miller PE, Levine JA, *et al.* Cost savings of reduced constipation rates attributed to increased dietary fiber intakes: a decision-analytic model. *BMC Public Health* 2014; **14**: 1–7.
64. Citronberg J, Kantor ED, Potter JD, White E. A prospective study of the effect of bowel movement frequency, constipation, and laxative use on colorectal cancer risk. *Am J Gastroenterol* 2014; **109**: 1640–9.
65. Cook IJ, Talley NJ, Benninga MA, Rao SS, Scott SM. Chronic constipation: overview and challenges. *Neurogastroenterol Motil* 2009; **21** (Suppl. 2): 1–8.