

Guido Camps PhD DVM

VoedingNL22 | Biologisch ritme en voeding | 17 juni 2022

KU LEUVEN

Europese Unie
Europees Fonds voor Regionale Ontwikkeling

WAGENINGEN
UNIVERSITY & RESEARCH

OnePlanet
Research Center



1

Biologisch ritme en techniek

Wat ga ik vertellen

Tracken van

- Stress
- Slaap
- Voeding



2

Biologisch ritme en techniek


Tracken van stress



3

Meten van hartslag

ECG is de gouden standaard




4



5

Photoplethysmography (PPG)




6



HUNGRY ROBOT

7

These results suggest changes in the morphology of PPG pulses possibly revealing marks of arterial stiffness occurring due to psychological stress.



Ref 2, 3

HUNGRY ROBOT

8

Biologisch ritme en techniek

Tracken van stress - conclusie

Therefore, the outcome of this study suggest the reliability of implementing PPG signal for stress recognition.

Gegeven de aanname dat hartslag gezien wordt als marker van stress, dan zijn deze metingen dus betrouwbaar genoeg om een indicatie te geven van stress.

HUNGRY ROBOT

9




HUNGRY ROBOT

10

Kleine inkijk in de toekomst

Apple seeks patent on technology that will be used to monitor Apple Watch users' blood sugar readings

If Apple can make the technology work, those with the Series 7 Apple Watch who are insulin dependent diabetics will be able to monitor their blood sugar without paying for the disposable test strips. More importantly, they won't have to feel the pain of pricking their finger with a tiny needle and wait a few seconds since many diabetics don't like to go through the process of obtaining a reading, the new feature could lead to improved monitoring on the part of these patients and help them control their diabetes better. And the Apple Watch will eventually pay for itself since there are no more test strips to buy.



11

Biologisch ritme en techniek


Tracken van slaap

HUNGRY ROBOT

12

Meten van slaap

polysomnography (PSG) is de gouden standaard



Ref 4

HUNGRY ROBOT

13

Wrist actigraphy is based on the principle that physical movements are increased during wakefulness and reduced during sleep




Ref 5, 6

HUNGRY ROBOT

14

Meten van slaap – wat is er op de markt?

software



Ref 4

HUNGRY ROBOT

15

Meten van slaap – wat is er op de markt?

Smart watch gebaseerd



Ref 4

HUNGRY ROBOT

16

Meten van slaap – wat is er op de markt?

Dedicated tracker




Ref 4

HUNGRY ROBOT

17

Meten van slaap

Validiteit



It has been found to have a reasonable degree of agreement with PSG, with reported agreement rates of 78.8–99.7% for sleep and 48.5–79.8% for wake

However, its validity in special populations such as the elderly, in subjects with poor sleep quality, or in those with major health problems is not well established

Ref 4

HUNGRY ROBOT

18

Biologisch ritme en techniek

Tracken van slaap - conclusie

Slaap kan gemeten worden, maar algoritmes geven geen zekerheid over hun resultaat.

Vooral bij de voor ons interessante subgroepen is er twijfel over de validiteit, maar het geeft wel een indicatie.



19

Biologisch ritme en techniek

Tracken van voeding



20

Kijkje in ons huidig onderzoek

Wanneer eten we?

Hoe eten we?

Wat eten we?



21

Wanneer eten we?



22




23

When do people eat?


Inbetween meal consumption




24




25




26



27



28




29




30

Eten

- Sensor: Shimmer3 IMU wristband
- Collected Dataset:
 - 24 meal sessions from 24 subjects
 - 1461 bites in total
- Model:
 - 1D-TCN
 - Conv1d
 - Dilated layer
 - Residual connection
- Results: F1-score 94.4%



31

Model: Multi-Stage Temporal Convolutional Network (MS-TCN)


32

Result : Leave-one-subject-out (LOSO) Method

33

Drinking**Challenges**


- Distribute in all over the day
- Sparse
- Non-specific hand



34

Data collection

- Dataset description:**
 - Dataset I (short term)- (DX-I)**
 - 13 participants,
 - 486 drinking events: 101(L):266(R):43(L&R)
 - 10.3 hour
 - Duration ratio left hand null : drinking = 32.7 : 1
 - Duration ratio right hand null : drinking = 21.6 : 1
 - Dataset II (long term)- (DX-II)**
 - 7 participants (from 10am to 4 or 5 pm)
 - 314 drinking events: 142 (L) :152 (R) : 10(L&R)
 - 45.2 hour
 - Duration ratio left hand null : drinking = 218.7 : 1
 - Duration ratio right hand null : drinking = 175.1 : 1
- Results: F1-score : 90.0%




35

3. Radar-based Eating Gesture Detection

- Sensor: TI FMCW radar**
- Collected Dataset:**
 - 22 meal sessions from 22 subjects
 - 1138 bites in total
- Input data:**
 - Range-Doppler Cube
 - 25 Frames per Second
 - 64 x 64 per frame
- Model:**
 - 3D-TCN
 - Conv1d
 - Dilated layer
 - Residual connection
- Results: F1-score 93.3%**



36



37



38



39




40



41




42




43




44



45



46



47



48

- Model parameters:** 3 hidden layers, adam solver, relu activation, 1000 epochs
- Evaluation metrics problem:**
 - multi label multi output classification, e.g. no confusion matrix or other straightforward metrics
- Custom accuracy score calculated;**
0.8146
- Conclusion:**
 - Higher granularity needed
→ macronutrients

HUNGRY ROBOT

49

Als dit allemaal werkt, dan weten we:

Wat je eet, wanneer je eet en hoe je eet, en dat geheel automatisch.

50

Biologisch ritme en techniek

Tracken van voeding - conclusie

Er komen mooie methoden aan, maar de grote vraag is of deze overeind blijven in de echte wereld. Voor de komende jaren blijven de 'ouderwets' methoden de belangrijkste bron van informatie.

HUNGRY ROBOT

51

Biologisch ritme en techniek

Uiteindelijke conclusie

Qua hartslag, stress en slap is een (relatief goedkope) tracker wellicht een slimme investering.

Qua voeding zijn we nog niet zo ver.

52

@GuidoCamps

@<https://www.linkedin.com/in/guidocamps/>

HUNGRY ROBOT

53

Referenties

1. Dzedzicki, Andrius & Kalkauskas, Arturas & Bučinskas, Vytautas. (2020). Human Emotion Recognition: Review of Sensors and Methods. *Sensors*, 20, 592.
2. 10.3390/20030592.
3. Banerjee, R. Ballón, J. Lázaro, V. Marcais, P. Laguna and E. Gil, "A two step Gaussian modelling to assess PNG morphological variability induced by psychological distress," *2017 IEEE/CIBEC*, 2017, 270–275.
4. Alami, Noor & Soekti, Khan & Marnoto, H. (2017). Stress Recognition Using Photoplethysmogram Signal. *Indonesian Journal of Electrical Engineering and Computer Sciences*, 8, 495-501. 10.11591/ieeccv.v8.i2.pp495-501.
5. Bhana Prakash Kolia, Subir Mansukhani & Meghn P. Mansukhani (2016) Consumer sleep tracking devices: a review of mechanisms, validity and utility, *Expert Review of Medical Devices*, 13, 5, 497-506, DOI: 10.1586/17434440.2016.1171708
6. Tilmanne J, Urban J, Kothare MV, et al. Algorithms for sleep-wake identification using actigraphy: a comparative study and new results. *J Sleep Res*. 2009;18(1):85–98.
7. Sadeh A. The role and validity of actigraphy in sleep medicine: an update. *Sleep Med Rev* 2011;15(4):259–267.

54